The Bryologist 93(4), 1990, pp. 531–538
Copyright © 1990 by the American Bryological and Lichenological Society, Inc.

Index to Volume 93 of THE BRYOLOGIST[1,2]

Abrothallus oxysporus, *211*
Acarospora chlorophana, 202; fuscata, 202; schleicheri, 202; sinopica, 358; smaragdula, 202
Acer rubrum, saccharum, epiphytic lichens and bryophytes on, 257
Acrocladium auriculatum, 401; chlamydophyllum, 401
Acrocordia cavata, 359
Aerobryopsis subdivergens, 400
Agrestia hispida, 202
Agyrium, 211; rufum, 211
AKIYAMA, H. Morphology and taxonomic significance of dormant branch primordia, dormant buds, and vegetative reproductive organs in the suborders Leucodontineae and Neckerineae (Musci, Isobryales), 395
Alabama, Fissidens neonii new to, and additional reports for F. asplenioides, F. fontanus, and F. grandifrons, 30.
ALBRIGHT, D. L. *See* SHAW, A. J. & D. L. ALBRIGHT, 187
AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN. Lipids of the gametophyte and sporophyte of Funaria hygrometrica. Comparison with lipids from leaves of vascular plants, 44
ALI, A. M. *See* AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN, 44
ALLEN, B. Campylopus atlanticus (Musci: Dicranaceae), a new species from Panama, with comments on the taxonomy of Campylopus, 445
ALLEN, B. H. A reconsideration of Dicranoweisia calcarea, 206
ALLEN, B. H. [Review of:] N. A. Konstantinova, O. A. Belkina, A. J. Lihacev & R. N. Sljkov (Eds.). Bryophyta Murmanica Exsiccata (USSR). Fasciculus I (No. 1–50), 100
ALLEN, B. H. [Review of:] S. W. Greene & A. J. Harrington. The conspectus of bryological taxonomic literature. Part 1. Index to monographs and regional reviews, 381
ALLEN, B. H. *See* BOWERS, F. D. & B. H. ALLEN, 346
ALLEN, B. H. *See* MAGILL, R. E. & B. H. ALLEN, 84, 368
ALLEN, B. H. & R. E. MAGILL. Recent literature on mosses—144, 242; —146, 520
Alsia, 396; californica, 398; longipes, *328*
Amblystegium serpens, 266; varium, 266
Amphiloma lanuginosum, *211+*
Anaptychia isidiza, 138; palmulata, 138, 264, 359; stippaea, 138
ANDERSON, L. E. A checklist of Sphagnum in North America north of Mexico, 500
ANDERSON, L. E., H. A. CRUM & W. R. BUCK. List of the mosses of North America north of Mexico, 448
Aneura pinguis, 62
Anisomeridium biforme, 359; nyssaegenum, 264
Anomodon abbreviatus, 398; attenuatus, 266; giraldii, 398; longifolius, 398; minor, 398; rostratus, 266+; rugelii, 266, 398; thraustus, 398
Anomodontaceae, 395+
Anthracothecium corticatum, *211+*; falsarium, *211+*; leucostomum, *211+*; maculare, *211+*; mucosum, *211*; nanum, 212; ochraceoflavens, *212+*; ochraceoflavum, *212+*; pauciloculare, *212*; pyrenuloides, *212*; staurosporum, 212; subglobosum, 212; thelomorphum, *212+*; varians, 212
Antitrichia californica, 398; curtipendula, 398
Anzia colpodes, 264; colpota, 109; opuntiella, 109
Arthonia arthonioides, 212+; aspersa, *212*; byssacea, 264, 359; diffusella, 359; fuliginosa, 212; leucopellaea, 212; patellulata, 359; phaeobaea, 212; rubella, 264; spadicea, 264
Arthothelium ruanum, 264; spectabile, 264
ARTS, T. & J.-P. FRAHM. Campylopus pyriformis new to North America, 290
Aspicilia alphoplaca, 202; caesiocinerea 202; candida 202; cinerea, 202
Aulocomnium palustre, 17, 63, 349

Bacidia accedens, *212+*; atrogrisea, 359; bacillifera, 359; bagliettoana, 359; carneoalbida, 212; chlorantha, 264; circumspecta, 359; epixanthoides, *360*; fusca, *212+*; globulosa, 212; granosa, 360; incompta, 359; laurocerasi, 359; microcarpa, *212+*; microphyllina, *212+*; naegelii, 359; obscurata, *212+*; polychroa, 264; populorum, 360; schweinitzii, 264; sphaeroides, *212*; tetramera, *212*; trachona, 360
Bactrospora patellarioides, 212+
Barbella asperifolia, 400; enervis, 400; pendula, 400
Barbula jacksharpii, *207*
Bazzania trilobata, 266+
Bellemerea alpina, 202
BELNAP, J. & K. T. HARPER. Effects of a coal-fired power plant on the rock lichen Rhizoplaca melanophthalma: Chlorophyll degradation and electrolyte leakage, 309
Belonia fennica, 212
BENEDICT, J. B. Winter frost injury to lichens—Colorado Front Range, 423
BENEDICT, J. B. & T. H. NASH III. Radial growth and habitat selection by morphologically similar chemotypes of Xanthoparmelia, 319
Bescherellia brevifolia, 399+*
Bestia, 328+; holzingeri, *330*; longipes, 328; occidentalis, *330*; vancouveriensis, *330*
Betula alleghaniensis, epiphytic lichens and bryophytes on, 257
Biatora botryosa, 212+; campestris, 364; epixanthoides, 360; helvola, 212+; sphaeroides, 212+
Biatorella albidula, *212+*; resinae, *212+*
Bissetia lingulata, 401
Blepharoplast morphology, Treubia tasmanica, 409
BLEWITT, M. R. & G. A. COOPER-DRIVER. The effects of lichen extracts on feeding by gypsy moths (Lymantria dispar), 220
Bolivia, Sphagnum inretortum, new species from, 283
Borneo, mosses new to, 429+
Botrydina, *212+*; botryoides, *212+*; luteovitellina, *212+*; velutina, *212+*; viridis, *212+*; vulgaris, *212+*
BOUCHER, V. L. & T. H. NASH III. Growth patterns in Ramalina menziesii in California: Coastal vs. inland populations, 295
BOWERS, F. D. & B. H. ALLEN. A new species of Brachydontium (Musci: Seligeriaceae) from Honduras, 346

[1] Compiled by Sharon Eversman, Biology Department, Montana State University, Bozeman, MT 59717-0346.

[2] New scientific names and combinations are in boldface; page numbers of names in synonomy are in italic; taxa mentioned several times in a paper are indicated by a plus sign (+) after the page number; illustrations are indicated by an asterisk (*).

0007-2745/90/531–538$0.95/0

BOWERS, F. D. & S. L. TIMME. Fissidens neonii new to Alabama, and additional reports for F. asplenioides, F. fontanus, and F. grandifrons for Alabama, 30
Brachydontium **olsonii** sp. nov., 346*
Brachythecium acuminatum, 266; curtum, 266; oxycladon, 266; rutabulum, 63; salebrosum, 266; velutinum, 266
Braithwaitea sulcata, 400
BRAKO, L. [Review of:] M. Galun. CRC Handbook of Lichenology, Volume 1, 379
Braunia alopecura, 398; attenuata, 398; secunda, 398; squarrulosa, 398
BRODO, I. M. [Review of:] T. L. Esslinger. Systematics of Oropogon (Alectoriaceae) in the New World, 255
BRODO, I. M. *See* WONG, P. Y. & I. M. BRODO, 357
Brotherella recurvans, 266+
BRUGGEMAN-NANNENGA, M. A. & R. A. PURSELL. The Fissidens radicans complex (Section Amblyothallia) in the neotropics and paleotropics, 332
Bryoerythrophyllum ferruginascens in eastern North America, 208*; inaequalifolium, 108; recurvirostrum, 208, var. recurvirostrum, 386*, var. robustum, 386*
Bryolawtonia gen. nov., 328+; **vancouveriensis** comb. nov., 329+
Bryoria abbreviata, 202; fremontii, 202; friabilis, 212; fuscescens, 202; implexa, 212; lanestris, 202; pseudofuscescens, 212; vrangiana, *212*
Bryowijkia ambigua, 398
Bryum argenteum, heavy metal tolerance in, 187; bicolor, 1+; incrassatolimbatum, 286+*; pseudotriquetrum, 63, 349; **tuberosum** sp. nov., 288*
BUCK, W. R. [Review of:] S. W. Greene & A. J. Harrington. The Conspectus of Bryological Taxonomic Literature. Part 2. Guide to National and Regional Literature, 98
BUCK, W. R. [Review of:] W. Frey (ed.). Mooseflora und -vegetation in Regenwaldern NO-Perus. Ergenbnisse der BRYOTROP-Expedition nach Peru 1982, 98
BUCK, W. R. *See* ANDERSON, L. E., H. A. CRUM & W. R. BUCK, 448
BUCK, W. R. *See* ZOMLEFER, W. B. & W. R. BUCK, 303
Buellia amphidexia, 212; arnoldii, 212, 264, 360; bahiana, 212; callispora, *212+*; catasema, 212; coccinea, 212; curtisii, 264; dialyta, 264+; disciformis, 264; elizae, 264; erubescens, 202; melanochlora, 212; modesta, 212; pachnidisca, 212; papillata, 202; placodiomorpha, 212; polyspora, 212, 360; rinodinaspora, *212+*; rubifaciens, 212; stillingiana, 67, 264; wheeleri, 212
BURKHALTER, J. R. *See* WILHELM, G. S. & J. R. BURKHALTER, 66

Calicium adspersum, 212; corynellum, 212; hyperelloides, 212; lenticulare, 212; minutissimum, *212*; quercinum, 212; roscidum, *212+*; subquercinum, *212+*; trabinellum, 212+
California, Ramalina menziesii growth patterns in, 295
Calliergon trifarium, 349
Calliergonella cuspidata, 63+
Caloplaca brattiae, 212; chrysophthalma, 360; citrina, 202, 360; decipiens, 202; discolor, 360; epithallina, 202; ferruginea, 202, 360; flavovirescens, 202, 360; fraudans, 202; microphyllina, 360; sideritis, 360; tiroliense, 202; trachyphylla, 202; velana, 360
Calymperes othmeri, 37*
Calypogeia, 314+; azurea, 314; fissa, 314, subsp. neogaea, 315; integristipula, 315; muelleriana, 316, fo. **schofieldii** fo. nov., 316, fo. **shieldsii** fo. nov., 316; neesiana, 316; sphagnicola, 317; suecica, 317
Calypogeiaceae, western North America, 313
Calyptothecium caudatum, 399; hookeri, 399; tumidum, 399
CAMERON, R. G. & R. WYATT. Spatial patterns and sex ratios in dioecious and monoecious mosses of the genus Splachnum, 161
Camptochaete porotrichoides, 401
Campylium hispidulum, 266; stellatum, 17+, 64
Campylopus, 445+; **atlanticus** sp. nov., 445*; pyriformis, 290+*
Canada, habitat limitations of Sphagnum in western, 7+
Candelaria concolor, 109+, 202, 264
Candelariella aurella, 202; vitellina, 202; xanthostigma, 202
Canoparmelia aptata, 125; caroliniana, 264
CAROTHERS, Z. B. & A. E. RUSHING. Blepharoplast morphology in Treubia tasmanica (Hepaticae: Treubiales), 409
Catapyrenium cinereum, 212; daedaleum, 212; globosum, 212; lachneum, 202; schaereri, 212; waltheri, *212*
Catascopium nigritum, 62, 350
Catillaria globulosa, *213+*; sphaeroides, *213+*
Catinaria laureri, 264
Cecidonia, 213; umbonella, 213
Cercidospora ulothii, 213
Cetrariastrum cirrhatum, 123
Cetrariopsis wallichiana, 122+
Cetrelia braunsiana, 123; cetrarioides, 264; chicitae, 123, 264; japonica, 123+; monachorum, 124; olivetorum, 124, 264; pseudolivetorum, 124
Chaenotheca brachypoda, 213+; carthusiae, *213*; chlorella, 213; gracilenta, 213
Chaenothecopsis savonica, 360
Checklist of mosses of North America north of Mexico, 448
Checklist of Sphagnum of North America north of Mexico, 500
chemistry, Ramalina americana complex, 167
chemotypes, Ramalina americana complex, 167
Chrysopsora testacea, *213+*
Chrysothrix candelaris, 264; chlorina, 202
CHURCHILL, S. P. Pseudocrossidium steerei (Pottiaceae), a new species from Ecuador, 353
Cinclidium stygium, 350
Ciadidium bolanderi, 212+; thamnitis, *213+*
Cladina, 213+; arbuscula, 148, evansii, 66; mitis, 149, 202, 213; pseudoevansii, 149; rangiferina, 66, 149; subtenuis, 66, 220+
Cladomnion ericoides, 399
Cladonia amarocraea, 150; bacillaris, 150; beaumontii, 66; caespitica, 150; cariosa, 202; carneola, 202; cenotea, 202; chlorophaea, 151, 202, 264; coniocraea, 151, 202, 264; cornuta, 264; cryptochlorophaea, 151; deformis, 202; didyma, 151; dimorphoclada, 360; ecmocyna, 202; fimbriata, 151, 202; floerkeana, 152; furcata, 152, 264; gracilis subsp. nigripes, 152, subsp. turbinata, 152; homosekikaica, 153; krempelhuberi var. sublepidota, 153; leporina, 66; magyarica, 213; mateochatha, 202; ochrochlora, 153; pachycladodes, 66; perforata, 66+; phyllophora, 153; pityrea, *213+*; pleurota, 153; prostrata, 66; pyxidata, 153, 202; ramulosa, 154; rappii var. exilior, 67; ravenelii, 67; scabriuscula, 154, 213; squamosa, 154, 264; subradiata, 67; subsquamosa, 213; sulphurina, 202; thiersii, 213; turgida, 154
Climacium americanum, 63
Coccocarpia erythroxyli, 110; palmicola, 110, 264; pellita, 110
Coelocaulon aculeatum, 202; muricatum, 202
Collema coccophorum, 202; flaccidum, 110; furfuraceum,

111; fuscovirens, 202; japonicum, 111; nigrescens, 111, 264; pulchellum var. subnigrescens, 213; shiroumanum, 111; subflaccidum, 111, 202, 264; undulatum, 202
Cololejeunea biddlecomiae, 266+
Colorado, winter damage to lichens in, 423; growth of Xanthoparmelia chemotypes in, 319
Coniocybe sulphurea, *213+*
Conomitrium radicans, *333*
COOPER-DRIVER, G. A. *See* BLEWITT, M. R. & G. A. COOPER-DRIVER, 220
Coriscium viride, *213+*
COXSON, D. S. & R. L. MACKEY. Diel periodicity of photosynthetic response in the subalpine moss Pohlia wahlenbergii, 417
CRANDALL-STOTLER, B. *See* STOTLER, R. E. & B. CRANDALL-STOTLER, 69, 222, 502
CRUM, H. Sphagnum inretortum, a new species in a new section from Bolivia, 283
CRUM, H. A. *See* ANDERSON, L. E., H. A. CRUM & W. R. BUCK, 448
Cryphaeaceae, 395+
Cryphaea nervosa, 398; obovatocarpa, 398, patens, 398; pilifera, 398; sinensis, 398
Cryphaeophilum, 396
Cryptogonium phyllogonioides, 397+*
Cryptolechia carneolutea, 361
Cryptoleptodon flexosus, 401
CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON. A novel *meta*-depside from the Ramalina subfraxinea complex, 193
CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON. The Ramalina americana complex (Ascomycotina, Ramalinaceae): Chemical and geographic correlations, 167
CULBERSON, C. F., A. JOHNSON, P. G. PATWARDHAN & U. MAKHIJA. New depsides in Stirtonia ramosa (Ascomycotina, Arthoniaceae), 279
CULBERSON, W. L. *See* CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON, 167, 193
Cyphelium notarisii, 361; tigillare, 202; trachylioides, 213
Cyptodon crassinervis, 398
Cyptodontopsis obtusifolia, 398
Cyrtopodaceae, 395+
Cyrtopus setosus, 399+*

Dactylina madreporiformis, 202
DAMANHURI, A. *See* MOHAMED, H. & A. DAMANHURI, 288
death notices, 102, 256
4-*O*-demethylsuperconfluentic acid, 279+
Dendriscocaulon umhausense, 264
Dendroalsia, 396; abietinea, 398; longipes, *328*
Dendrocryphaea gorveana, 398
Dendropogonella rufescens, 398
meta-depside, of Ramalina subfraxinea, 193
depsides, two new in Stirtonia ramosa, 279+
Dermatocarpon miniatum, 108, 202; myogiense, 108; vagans, 202; waltheri, *213+*
Diaphanodon blandus, 399; procumbens, 399
Dichelodontium nitidum, 399
Dicranella calcarea, *207*
Dicranoweisia calcarea, *206*
Dicranum montanum, 266+; scoparium, 266; viride, 266+
Dimelaena oreina, 202, 425
Dimerella lutea, 264
Diphyscium **buckii** sp. nov., 429*
Diploschistes scruposus, 203
Diplotomma epipolium, 361
Dirinaria aegialita, 213
Distichophyllum **noguchianum** sp. nov., 429*
Dolichomitra cymbifolia var. subintegerrima, 401
Dozya japonica, 398
Drepanocladus aduncus var. polycarpus, 63; crassicostatus, 351; lapponicus, 351; lycopodioides, 351; revolvens, 62, 349; sendtneri, 62, 351; uncinatus, 266; vernicosus, 62, 349
Drummondia prorepens, 266+
Duthiella flaccida, 399; speciosissima, 399; wallichii, 399

Echinodiaceae, 395+
Echinodium hispidum, 401
ECKEL, P. M. Bryoerythrophyllum ferruginascens (Musci: Pottiaceae) in eastern North America, 208
ECKEL, P. M. Taxa minora delineata. II, 386
Ecuador, Pseudocrossidium steerei new to, 353
EGAN, R. S. Changes to the "Fifth checklist of the lichen-forming, lichenicolous and allied fungi of the continental United States and Canada." Edition II, 211
EGAN, R. S. Recent literature on lichens—138, 76;—139, 229;—140, 508
EL-SAADAWI, W. E. *See* AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN, 44
Endocarpon pulvinatum, 203
ENROTH, J. *See* NORRIS, D. H. & J. ENROTH, 328
enzymes, Umbilicaria vellea, 389+
Eopyrenula leucoplaca, 264
Ephebe lanata, 361; solida, 203
epiphyte abundance, estimation of, 39+
Eumyurium sinicum, 399
Euptychium setigerum, 400
Eurhynchium hians, 63; praelongum, 303+
Evernia mesomorpha, 203, 264; prunastri, 32+
EVERSMAN, S. Lichens of Yellowstone National Park, 197

Fagus grandifolia, epiphytic lichens and bryophytes on, 257
FAHSELT, D. *See* HAGEMAN, C. & D. FAHSELT, 389
Felipponea esquiroli, 398; montevidense, 398
FIFE, A. J. [Review of:] A. Eddy, A handbook of Malesian mosses. Volume 1. Sphagnales to Dicranales, 97
Fissidens **allenianus** sp. nov., 332*; asplenioides, 30+; **dendrophilus** sp. nov., 335*; fontanus, 31; grandifrons, 31; guineensis, *339*; konkourae, *339*; microcarpus, 339*; neonii, 30; radicans, 333*; subsphaerocarpa, *339*
Flavoparmelia caperata, 125, 220+, 264, 427
Flavopunctelia flaventior, 428; soredica, 428
FLETCHER, A. *See* HALE, M. E. & A. FLETCHER, 23
Floribundaria aurea subsp. nipponica, 400; floribunda, 400
Forssiroemia, 396; cryphaeoides, 398; japonica, 398; neckeroides, 398; trichomitria, 398
fossil records, moss, 351
FRAHM, J.-P. *See* ARTS, T. & J.-P. FRAHM, 290
Frullania asagrayana, 266+; brittoniae, 267; eboracensis, 267+
Fulgensia bracteata, 203; fulgens, 203
Funaria hygrometrica, 44+

Gammiella **touwii** sp. nov., 432*
Garovagliaceae, 395+
Garovaglia angustifolia, 400, fo. bogoriensis, 400; baeuerlenii, 400; compressa, 400; elegans, 397*, fo. elegans, 400, subsp. dietrichiae, 400
GIGNAC, L. D. & D. H. VITT. Habitat limitations of Sphagnum along climatic, chemical, and physical gradients in mires of western Canada, 7
Graphina acrophaea, *213+*; parilis, 213+
Graphis scripta, 264+
Greenland, Meesia triquetra in, 349
growth patterns, Ramalina menziesii, 295

Gyalecta truncigena, 361
Gyalideopsis athalloides, 213; vainioi, 213
Gymnostomum **jacksharpii** comb. nov., 206
gypsy moths, effects of lichen extracts on feeding, 220

Haematomma cismonicum, *213+*; elatinum, *213+*; ochrophaeum, *213+*; pustulatum, *213+*, 264
HAGEMAN, C. & D. FAHSELT. Multiple enzyme forms as indicators of functional sexuality in the lichen Umbilicaria vellea, 389
Hageniella **hattoriana** sp. nov., 433*
HALE, M. E. & A. FLETCHER. Rimelia Hale & Fletcher, a new lichen genus (Ascomycotina: Parmeliaceae), 23
Hampeella pallens, 399
Handeliobryum sikkimense, 401
Haplohymenium pseudo-triste, 398; sieboldii, 398; triste, 398
HARPER, K. T. *See* BELNAP, J. & K. T. HARPER, 309
Hedwigia, 396; ciliata, 398+*
Hedwigiaceae, 395+
Hedwigidium integrifolium, 398
Helicodontium complanatum, 308; pervirens, 307
Henicodium, 396; geniculatum, 397+*
Heppia euploca, *213+*; guepinii, *213+*; polyphylla, *213+*
Herpetineuron toccoae, 398
Heterocladium vancouveriense, *329*
Heterodermia boryi 139; dendritica, 139; diademata, 139; firmula, 140; hypoleuca, 140; isidiophora, 140; japonica, 140; leucomelos, 264; cf. loriformis, 141; microphylla, 141; obscurata, 264; propagulifera, 141; pseudospeciosa, 141; speciosa, 141, 264; squamulosa, 264
Homalia japonica, 401; trichomanoides, 401
Homaliadelphus targionianus, 401+*
Homaliodendron exiguum, 401; scalpellifolium, 401
Homomallium adnatum, 266
Honduras, Brachydontium olsonii, new species from, 346
HONG, W. S. The family Calypogeiaceae in North America west of the hundredth meridian, 313
Horikawaea nitida, 397+*
HORTON, D. G. *See* VITT, D. H. & D. G. HORTON, 62
Huilia nigrocruenta, *213+*
Hygrohypnum reduncum, 278+
Hylocomium splendens, 17, 266
Hymenelia ceracea, 213; lacustris, 361
Hyperphyscia adglutinata, *213+*; minor, *213+*
Hypnum lindbergii, 63; occidentale, *330*; pallescens, 266+; pterigynandroides, *307*
Hypocenomyce scalaris, 203
Hypogymnia austerodes, 203; cf. enteromorpha, 137; fragillima, 137; imshaugii, 203; oceanica, 213; physodes, 203, 220+, 264; pseudophysodes, 137; pulverata, 213
Hypotrachyna croceopustulata, 264+; imbricatula, 264+; osseoalba, 128; revoluta, 264; rockii, 264

Idaho, lichens as air pollution biomonitors in, 50
Indonesia, mosses new to, 492+
Ionaspis, 213; arctica, 213+; epulotica, 213+, 361, var. arctica, *213+*; odora, 213; suaveolens, 213
IRELAND, R. R. Isopterygium acutifolium, a new moss species from Venezuela, 343
Isopterygium **acutifolium** sp. nov., 343*; affusum, 343; aquaticum, 434; byssobolax, 343; herminieri, 343; irelandii, 343; jamaicense, 343; subbrevisetum, 343; subglobosum, 343; submersum, 343; tenerifolium, 343; tenerum, 343
Isothecium holzingeri, *330*; occidentale, *330*

Jaegerina latiscuspis, 399; scariosa, 399
Jamesoniella autumnalis, 267
JOHNSON, A. *See* CULBERSON, C. F., W. L. CULBERSON & A. JOHNSON, 167, 193
JOHNSON, A. *See* CULBERSON, C. F., A. JOHNSON, P. G. PATWARDHAN & U. MAKHIJA, 279

KANDA, H. *See* NISHIMURA, N. & H. KANDA, 275

Lecanactis, 213; chloroconia, 264, 361; grumulosa, *213+*; patellarioides, *213+*
Lecania albariella, *213+*; turicensis, 213+
Lecanora actophila, 213; allophana, 264; argentata, 264; argopholis, 203; atriseda, *213+*; bolanderi, *213+*; caesiorubella, 264; cenisia, 203; chloroconia, 361; chlorophaeodes, 213, 361; cinereofusca, 264+; crenulata, 203, 361; epulotica, *213+*; garovaglii, 203; glabrata, 264; hybocarpa, 264; hypospilota, *213+*; impudens, 264; imshaugii, 264; melanophthalma, 52+; muralis, 203, 428; mutabilis, *213+*; nigromarginata, 203; novomexicana, 203; ochrococca, *213+*; pallida var. rubescens, 265; phaeobola, *213+*; piniperda, 203; polytropa, 203; rugosella, 265; rupicola, 203; saligna, 203; strobilina, 265; subrugosum, 265; tesselina, *213+*; thamnitis, *213+*; thomsonii, 203; thysanophora, 265; umbrina, 213; urceolata, *213+*; verrucosa, *213+*
Lecidea atrobrunnea, 203, 425; atrofusca, *213+*; berengeriana, 203, *213+*; botryosa, *214+*; contigua, *214+*; cyrtidia, 361; diversa, *214+*; elabens, 361; erythrophaea, 265, 362; fusca, *214+*; helvola, *214+*, 265; hypnorum, *214+*; lactea, 214+; ochrococca, *214+*; oreinodes, 214+; oxyspora, *214+*; phylliscina, *214+*; plebeja, 362; sanguineoatra, *214+*; templetonii, *214+*; tennessensis, *214+*; tessellata, 203; tesselina, *214+*
Lecidella elaeochroma, 362; euphorea, 203, 362; stigmatea, 203
Lembophyllaceae, 395+
Lembophyllum bolivianum, 401; clandestinum, 401
Lepidozia reptans, 267+
Lepraria arctica, *214+*; incana, 203; membranacea, 203, *214+*; neglecta, 203; vouauxii, *214+*
Leprocaulon albicans, 203
Leproloma, 214; angardianum, 214; diffusum var. diffusum, 214; membranaceum, *214+*; vouauxii, *214+*
Leptodon smithii, 401+*
Leptogium azureum, 112; corticola, 265; crenatellum, *214+*; cyanescens, 112, 265; hirsutum, 112; menziesii, 112; moluccanum, 113; rivulare, 214+, 362; saturninum, 113, 203; tenuissimum, 203; cf. teretiusculum, 362
Lepyrodon australis, 399; lagurus, 399; tomentosus, 399
Lepyrodontaceae, 395+
Leskeella nervosa, 266+
Letharia columbiana, 203; vulpina, 203
Leucodon brachypus, 266+; pendulus, 398; sciuroides, 398; spp. 398
Leucodontaceae, 395+
Leucodontineae, 395+
Leucolejeunea clypeata, 267
Lichen perlatus α ciliatus, *28*
Lichenochora, 214; weillii, 214
Lichenosticta, 214; alcicornaria, 214
lichens affected by coal-fired power plant, 309; as air pollution biomonitors, 50; and gypsy moth larvae, 220; in Yellowstone National Park, 197; recolonization of, 427
Limprichtia **cossonii** comb. nov., 463
lipids of Funaria hygrometrica, 44+
List of mosses of North America north of Mexico, 448; of Sphagnum of North America north of Mexico, 500
Lithographa, 214; tesserata, 214

Lobaria adscripturiens, 118; japonica, 119; linita, 119; meridionalis, 119; pulmonaria, 265; quercizans, 119; retigera, 120; spathulata, 120
LOUN, J. Thamnolia vermicularis, 104
Loxospora cismonica, 214+; elatina, 214; ochrophaea, 214; pustulata, 214

MACKEY, R. L. *See* COXSON, D. S. & R. L. MACKEY. 417
macrolichen flora of South Korea, 105
MAGILL, R. E. *See* ALLEN, B. H. & R. E. MAGILL, 242
MAGILL, R. E. & B. H. ALLEN. Recent literature on mosses—143, 84; —145, 368
MAKHIJA, U. *See* CULBERSON, C. F., A. JOHNSON, P. G. PATWARDHAN & U. MAKHIJA, 279
Malaysia, Bryum tuberosum new to, 288
Malesia, mosses new to, 429
MCCUNE, B. Rapid estimation of abundance of epiphytes on branches, 39
MCCUNE, B. *See* STONE, D. F. & B. MCCUNE, 32
Meesia triquetra, 17+, 62, 349+
Megaspora, 214; verrucosa, 214+
Melanelia exasperatula, 203; fuliginosa, 265; granulosa, 203; halei, 265; huei, 124; incolorata, 203; infumata, 203; olivacea, 125; sorediata, 203; stygia, 203; subaurifera, 265; subelegantula, 203; subolivacea, 203; substygia, 203, 424
Melanotheca, 214; achariana, *214+*; aggregata, *214*; anomala, *214+*; cinerata, *214+*
Menegazzia cf. asahinae, 137; terebrata, 137, 265
Mesoamerica, Ramalina americana complex from, 167+
Metacalypogeia, 313+; schusterana, 314
metals in Bryum argenteum, 187
Metamelaena, 214; melambola, 214+
Metaneckera menziesii, 401+*
Meteoriaceae, 395+
Meteoriella soluta, 400
Meteoriopsis reclinata, 400
Meteorium helminthocladum, 400; miquelianum, 400
2′-*O*-methylnorsuperphyllinic acid, 279+
4-O-methyloxocryptochlorophaeic acid, Ramalina subfraxinea complex, 193
Metzgeria crassipilis, 267
Micarea endocyanea, 265+; hedlundii, 214; lutulata, 214; peliocarpa, 362; prasina, 363
Microcalicium arenarium, 362
MISHLER, B. D. [Review of:] R. N. Chopra & P. K. Kumra. Biology of Bryophytes, 100
Missouri, fen bryophytes in, 62
Mittenia plumula var. **gigantea** var. nov., 433*; var. plumula, 435*
Mittenothamnium **deguchii** sp. nov., 275*; scalpellifolium, 278
Miyabea fruticella. 398
Moenkemeyera konkourae, *339*; subsphaerocarpa, *339*
MOHAMED, H. & A. DAMANHURI. Bryum tuberosum, a new species from peninsular Malaysia, 288
MONTAGNES, R. J. S. The habitat and distribution of Meesia triquetra in North America and Greenland, 349
mosses of North America north of Mexico, list of, 448
Mycobilimbia, 214; accedens, 214+; berengeriana, 214+; fusca, 214+; hypnorum, 214+; lobulata, 214+; microcarpa, 214+; obscurata, *214+*; ohiense, *214+*
Myelochroa aurulenta, 132, 428; **coreana** sp. nov., 132*; crassata, 132; entotheiochroa, 133; galbina, 134; hayachinensis, 135; indica, 134; irrugans, 134; rhytidodes, 135; subaurulenta, 135
Myrionora, 214; albidula, 214+; clandestinum, 214+; compunctum, 214; glaucophaeum, 214+; terebratulum, 214+
Myuriaceae, 395+

Nadvornikia, 214; hawaiensis, 214
NASH, T. H. III. *See* BENEDICT, J. B. & T. H. NASH III, 319
NASH, T. H. III. *See* BOUCHER, V. L. & T. H. NASH III, 295
Neckera borealis, 401; humilis, 401; pennata, 266+*, 401; yezoana, 401
Neckeraceae, 395+
Neckerineae, 395+
Neckeropsis calcicola, 401; gracilenta, 401; lepineana, 401; nitidula, 401; obtusata, 401
Neolindbergia brassii, 399; flagellifera, 399; plicata, 399; rigida, 399; robusta, 399; rugosa, 399
Nephroma helveticum, 113, 265; parile, 113, 203; resupinatum, 113; servitianum, 114
Nephromopsis asahinae, 121; ornata, 122; **pallescens** comb. nov., 122; pseudocomplicata, 122; stracheyi, 122
Nesolechia, *214*; oxyspora, *214+*
New Caledonia, chemistry of Ramalina subfraxinea, 193
New York, lichens and bryophytes in Adirondack Mountains, 257
NISHIMURA, N. & H. KANDA. Mittenothamnium deguchii, a new moss species from southern Chile, 275
Normandina pulchella, 108, 265
NORRIS, D. H. & J. ENROTH. Description of Bryolawtonia Norris & Enroth (Thamnobryaceae), with observations on the genus Bestia, 328
North America north of Mexico, list of mosses, 448; list of Sphagnum, 500
North Carolina, lichens and bryophytes in Blue Ridge Mountains of, 257

Ocellularia abrosella, *214+*; glaucophaea, *214+*; meiosperma, 214+; terebratula, *214+*
OCHI, H. & N. SALAZAR ALLEN. Bryum incrassatolimbatum new to Panama, with first description of its sporophyte, 286
Ochrolechia arborea, 265; pallescens, 67; pseudopallescens, 363; rhodoleuca, 214+; subplicans subsp. hultenii, 214+, subsp. subplicans, 214+; trochophora, 363; upsaliensis, 203
Oedicladium doii, 399; fragile, 399; rufescens, 399
Ohio River valley, lichen recolonization, 427
Omphalina, *214*; ericetorum, *215+*; hudsoniana, *215+*; luteovitellina, *215+*; velutina, *215+*
Ontario, lichen flora, 357
Opegrapha calcarea, 215; cinerea, *215+*; diaphoroides, *215+*; rimalis, *215+*; rupestris, 215+; saxicola, *215+*; varia, 215; viridis, 215; vulgata, *215+*
ORBÁN, S. & W. D. REESE. Syrrhopodon prolifer (Musci: Calymperaceae): A world view, 438
Oregon, annual branching in the lichen Evernia prunastri in, 32
Oropogon asiaticus, 147
Orthorrhynchium elegans, 399
Orthostichidium guyanense, 400
Orthostichopsis auricosta, 400; crinita, 400; perpinnatum, 400; subimbricata, 400; tenuis, 400
Orthotrichum sp., 266

Pachyospora, *215*; mutabilis, *215+*; verrucosa, 203, *215+*
Palisadula chrysophylla, 399+*; katoi, 399
Paludella squarrosa, 17+
Panama, Bryum incrassatolimbatum new to, 287; Campylopus atlanticus in, 445
Pannaria ahlneri, 114; conoplea, 114, 265; leucophaea, 203; leucosticta, 114; lurida, 115; praetermissa, 203; rubignosa, 115
Papillidiopsis **koponenii** sp. nov., 437*
Paraleucobryum longifolium, 266+

Parathelium, *215*; cuyabense, *215+*; emergens, *215+*; martinicanum, *215+*; microcarpum, *215+*; subferrugineum, *215+*
PARK, Y. S. The macrolichen flora of South Korea, 105
Parmelia sect. Hypotrachyna *Irregulares, *23*; sect. Irregulares, *23*; sect. Reticulatae, *23*; acanthifolia, *27*, fo. ornata, *28*; adaugescens, 126; amphygymnioides, *29*; angustata, *27*; badia, *27*; cetrata, *26*, fo. granularis, *29*, fo. platyloba, *29*, fo. sorediifera, *28*, subsp. radiata, *27*, var. corniculata, *27*, var. subisidiosa, *29*, var. ulcerata, *29*; ciliata, *29*; cinerascens var. saxicola, *27*; clavulifera, *29*; cochleata, 126; commensurata, *27*; concors, *28*; crinitum fo. varians, 363; decorata, *29*; diffractaica, *27*; diffusoides, *29*; exasperatula, 52+; fertilis, 126; foliolosa, *29*; ghattensis, *28*; hawaiiensis, *28*; herrei, *27*; homotoma, *28*, fo. incisocrenata, *28*, fo. subcervicornis, *28*; hypopsila, *215+*; ioanis-simae, *215+*; kurokowae, *215+*; laevigata var. reticulata, *28*; laevior, 127; livido-tessellata, *28*; macrocarpa, *28*; macrocarpoides, *28*, var. subcomparata, *29*; maura, *27*; menziesii, *27*; mutata fo. decorata, *28*; neoconspersa, *215+*; odontata var. rubiginosa, *29*; olivaria, *28*; ornata, *29*; owaniana, *27*; perforata fo. integra, *27*, var. corniculata, *27*, var. microspora, *27*, var. replicata, *27*; perlata α ciliatus, *28*, β olivaria, *28*, β olivetorum, *28*, g ciliata, *28*, var. ciliata, *28*, var. tentaculata, *28*; polita, *26*; praeperlata, *28*; pseudolaevior, 127; pseudoreticulata, *28*; ramescens, *28*; reterimulosa, *29*; reticulata, *28*, fo. nuda, *28*; ruminata, *28*; sanctae-helenae, *29*; saxatilis, 203; shinanoana, 128; simulans, *29*; squarrosa, 127, 265+; subisidiosa, *29*; sulcata, 203, 220+, 265, 428; taractica, *215+*; urceolata var. subcetrata, *28*; warmingii, *27*
Parmeliella cyanolepra, 203; incisa, 115; nigrocinerea, 115; subincisa, 115
Parmelina aurulenta, 265; galbina, 265; horrescens, 265; minarum, 265+; quercina, 135
Parmelinopsis expallida, 136; minarum, 136, 428; subfatiscens, 136
Parmeliopsis ambigua, 203; hyperopta, 203
Parmentaria, *215*; astroidea, *215+*; nana, *215+*; rappii, *215+*; ravenelii, *215+*
Parmotrema arnoldii, 129, 265; austrocetratum, *26*; cetratum, *27*, *265*; chinense, 129, 265; commensuratum, *27*; crinitum, 265+; diffractaicum, *27*; cf. eciliatum, 129; homotomum, *28*; hypotropum, 220+, 265; macrocarpum, *28*; margaritatum, 363; mellissii, 265; reticulatum, *29*, 129, 265+; robustum, 265+; simulans, *29*; stuppium, 265; subisidiosa, *29*, 265+; subtinctorium, 130, 363; tinctorum, 130
PATWARDHAN, P. G. *See* CULBERSON, C. F., A. JOHNSON, P. G. PATWARDHAN & U. MAKHIJA, 279
PEARSON, L. C. *See* ROPE, S. K. & L. C. PEARSON, 50
peat fens, 62+
Peltigera aphthosa, 203; canina, 116, 203; collina, 116, 203; degenii, 116; leucophlebia, 116; malacea, 117, 203; neopolydactyla, 117; nigripunctata, 117; polydactyla, 117, 203, 265; praetextata, 117; pruinosa, 118; rufescens, 203; scabrosa, 118; venosa, 203
Peltula euploca, 215+
Penzigiella cordata, 401
Pertusaria amara, 265; freyi, *215+*; globularis, 265; hultenii, *215+*; macounii, 265; multipunctoides, 265; ophthalmiza, 265; ostiolata, 265; paratuberculifera, 265; rhexostoma, 363; rhodoleuca, *215+*; rubefacta, 265+; subpertusa, 265; subplicans, *215+*; trachythallina, 265; velata, 265; waghornei, 265
Phacopsis, 215; oxyspora, 215+
Phaeophyscia adiastola, 142; endococcina, 203, 363; endococcinodes, 142; cf. erythrocardia, 142; exornatula, 143; hirsuta, 215+; cf. hirtella, 143; hirtuosa, 143; hispidula, 215+; imbricata, 143; limbata, 144; nepalensis, *215+*; cf. orbicularis, 144, 203, 265; primaria, 144; rubropulchra, 143, *215+*, 256+, 428; sciastra, 203
Phaeorrhiza nimbosa, 203
Phaeotrema meiospermum, *215+*
Philippines, mosses new to, 429+
Phlyctidia ludoviciensis, 265
Phlyctis 215; argena, 265
photosynthesis, Pohlia wahlenbergii, 417
Phyllogoniaceae, 395+
Phyllogonium fulgens, 401; immersus, 401
Phyllopsora corallina var. santensis, 215+; santensis, *215+*
Physcia adscendens, 203, 428; aipolia, 203; alba var. obsessa, *215+*; caesia, 203, 424; callosa, 203; dubia, 203; endochrysea, *215+*; hirsuta, *215+*; hispidula, *215+*; integra, 215+; millegrana, 265, 428; minor, *215+*; obsessa, *215+*; orbicularis fo. rubropulchra, *215+*; phaea, 203; setosa, *215+*; stellaris, 203; tenella subsp. marina, 215
Physciella denigrata, 144; nepalensis, 215+
Physciopsis minor, *215+*
Physcomitrium pyriforme, 63
Physconia detersa, 145, 203, 265; enteroxantha, 203, 363; grumosa, 145; kurokawae, 363; muscigena, 203; perisidiosa, 364
Phytoconis, 215; ericetorum, 215+; luteovitellina, 215+; velutina, 215+; viridis, 215+
Piloecium pseudorufescens, 399+*
Pilotrichella mauiensis, 401
Pilotrichopsis dentata, 398
Pinnatella makinoi, 401; sp. 401
Pireella cymbifolia, 400; formosana, 400; ludoviciae, 400; pohlii, 400
Placynthiella oligotropha, 364; uliginosa, 203
Placynthium nigrum, 203
Plagiocarpa hyalospora, 265+
Plagiomnium ciliare, 266; ellipticum, 63
Plagiothecium laetum, 266+
Platygyrium repens, 266+
Pleurotheliopsis australiensis, *215+*; nana, *215*
Pleurozium schreberi, 17+
Pohlia wahlenbergii, 417+
Polyblastia cf. obsoleta, 364
Polychidium muscicola, 203
Polycoccum, 215; epicrassum, *216+*; squamarioides, 216
Polypyrenula, *216*; sexlocularis, *216*
Polytrichum juniperinum, 17; **papillatum** comb. nov., 466
Porella platyphylloidea, 267+
Porothamnium bigelovii, 401; comosum, 401; pennaefrondeum, 401
Porotrichodendron superbum, 401
Porotrichum cobanense, 401; fruticosum, 401; plicatulum, 401; vancouveriense, *330*
Porpidia, 216; calcarea, 216; carlottiana, 216; contrapoenda, 216; diversa, 216+; flavicunda, 216; grisea, 216; herteliana, 216; lowiana, 216; macrocarpa, 216+; nigrocruenta, *216+*; tahawasiana, 216; thomsonii, 216
Prionodon ciliatus, 399; densus, 399; dichotomus, 399
Prionodontaceae, 395+
Protoparmelia atriseda, 216+; badia, 203; ochrococca, 216+
Pseudephebe pubescens, 204
Pseudobarbella attenuata, 401; levieri, 401
Pseudobraunia, 396; californica, 398
Pseudocrossidium steerei sp. nov., 353*
Pseudocryphaea domingensis, 396, 398
Pseudocyphellaria aurata, 265; crocata, 120
Pseudoleskea occidentalis, *329*; vancouveriensis, *329*
Pseudoleskeella occidentalis, *330*; vancouveriensis, *330*

Pseudoparmelia caperata, 269; caroliniana, 268
Pseudospiridentopsis horrida, 399
Psilolechia lucida, 364
Psora decipiens, 204; globifera, 204; nipponica, 204; russellii, 204
Psoroma hypnorum, 204
Pterobryaceae, 395+
Pterobryon arbuscula, 400+*; densum, 400; subarbusculum, 400
Pterobryopsis acuminata, 400; auriculata, 400+*; crassiuscula, 400; divergens, 400; flagellifera, 400; frondosa, 400; gedehensis, 400; mexicana, 400; orientalis var. gracilis, 400, var. yuennanensis, 400; schmidii, 400
Pterogonium, 396; gracile, 398
Ptilidium pulcherrimum, 267+
Ptychomniaceae, 395+
Ptychomnion aciculare, 399
Punctelia appalachensis, 266; borreri, 128, 266; reddenda, 266; rudecta, 128, 266+, 428; subrudecta, 266, 428
PURSELL, R. A. *See* BRUGGEMAN-NANNENGA, M. A. & R. A. PURSELL, 322
Pylaisiella intricata, 266; selwynii, 266
Pyrenastrum, *216*; cubanum, *216*+; fuscum, *216*+; pyrenastraeum, *216*+
Pyrenopsis melambola, *216*+; multispora, 216
Pyrenula, 216; acutalis, 216; anomala, 216+; aquila, 216; astroidea, 216; citriformis, 216; concatervans, *216*+; confoederata, 216; corticata, 216+; cubana, 216+; cuyabensis, 216+; erumpens, 216+; falsaria, 216+; furvella, 216; laevigata, 266; leucostoma, 216+; lucifera, 216; macularis, 216+; martinicana, 216+; micheneri, 216; microtheca, 216+; ochraceoflava, 216+; ochraceoflavens, 216+; oleosa, 216; plittii, 216, 266; pseudobufonia, 266+; punctelia, 266; pyrenuloides, 216+; ravenelii, 216; rubrostoma, 216; septicollaris, 216+; subaggregata, *216*; subelliptica, 266; subferruginea, 216+; tenuisepta, 216; thelomorpha, 216+
Pyrgillus americanus, *216*+; javanicus, 216+
Pyxine albovirens, 216+; caesiopruinosa, *216*+; endochrysina, 145; limbulata, 145; sorediata, 266

Racocarpus alpinus, 398; humboldtii, 398; purpurascens, 398
Racomitrium lanuginosum, 17+
Radula complanata, 267+
RADWAN, S. S. *See* AL-HASAN, R. H., W. E. EL-SAADAWI, A. M. ALI & S. S. RADWAN, 44
Ramalina americana, morphology of, 167; secondary chemistry of, 167; secondary compounds new to, 167; calicaris, 146; fastigiata, 167; menziesii, 295+; sayreana morph, 167+; siliquosa complex, 167+; sinensis, 147; subfraxinea complex, 193+; subgeniculata, 147; **unifolia** sp. nov., 341*; willeyi, 67; yasudae, 147
REESE, W. D. Calymperes othmeri reinstated, 37
REESE, W. D. *See* ORBÁN, S. & W. D. REESE, 438
Renauldia hoehnelii, 400; mexicana, 400
Rhizocarpon disporum, 204; geographicum, 204; lavatum, 364; macrosporum, 216+; oederi, 364; polycarpum, 362; riparium subsp. lindsayanum, 425; saanaense, 216+; sphaerosporum, *216*+; sublucidum, *216*+; submodestum, 216; subtile, *216*+; viridiatrum, 216
Rhizoplaca, chlorophyll degradation, electrolyte leakage of, 309; chrysoleuca, 204, 424; melanophthalma, 204, 425; peltata, 204
Rigodium, *303*+; arborescens, 401; crassicostatum, *306*; kilimandscharicum, 401; pterigynandroides, *307*+; riparium, *308*+; toxarioides, *303*+; toxarion, 401
Rimelia gen. nov., 23; **austrocetrata** comb. nov., 26; **cetrata** comb. nov., 26*; **commensurata** comb. nov., 27; **diffractaica** comb. nov., 27; **hawaiiensis** comb. nov., 28; **homotoma** comb. nov., 28; **macrocarpa** comb. nov., 28; **olivaria** comb. nov., 28; **reticulata** comb. nov., 28*; **ruminata** comb. nov., 38*; **simulans** comb. nov., 29*; **subsidiosa** comb. nov., 29
Rimularia insularis, 204
Rinodina archaea, 204; ascociscana, 266; bischoffii, 364; confragosa, 204; efflorescens, 364; farinosa, 364; gennarii, 204; olivaceobrunnea, 204; populicola, 364; subminuta, 266, 364; tephraspis, 364; thujae, 364
ROGERSON, C. T. [Review of:] V. Alstrup & D. L. Hawksworth. The lichenicolous fungi of Greenland, 378
ROPE, S. K. & L. C. PEARSON. Lichens as air pollution biomonitors in a semiarid environment in Idaho, 50
RUSHING, A. E. *See* CAROTHERS, Z. B. & A. E. RUSHING, 409
Rutenbergia armata, 399; limbata, 399; madagassa, 399; prionodon, 399
Rutenbergiaceae, 395+

SALAZAR ALLEN, N. *See* OCHI, H. & N. SALAZAR ALLEN, 286
Sarcosagium campestre, 364
Sarea, 216; difformis, 216; resinae, 216+
Schismatomma abietinum, 216+; pericleum, *216*+
Schistidium **hetrophyllum** comb. nov., 468
Schlotheimia brownii, 387*
SCHMITT, C. K. & N. G. SLACK. Host specificity of epiphytic lichens and bryophytes: A comparison of the Adirondack Mountains (New York) and the Southern Blue Ridge Mountains (North Carolina), 257
Schoenobryum coffeae, 398; concavifolia, 398; gardneri, 398
Scorpidium scorpioides, 62, 349; turgescens, 62
secondary products, new, Ramalina americana, 167+; Stirtonia ramosa, 279
SELVA, S. B. [Review of:] M. Galun (ed.). Handbook of lichenology. Volume II, 380
Sematophyllum adnatum, 266+
sex ratios, Splachnum, 161
SHAW, A. J. Genetic and environmental effects on morphology and asexual reproduction in the moss, Bryum bicolor, 1
SHAW, A. J. & D. L. ALBRIGHT. Potential for the evolution of heavy metal tolerance in Bryum argenteum, a moss. II. Generalized tolerance among diverse populations, 187
SHOWMAN, R. E. Lichen recolonization in the upper Ohio River valley, 427
Skytella, 216; mulleri, 216
SLACK, N. G. *See* SCHMITT, C. K. & N. G. SLACK, 257
Solorina crocea, 204
South Korea, macrolichen flora of, 105
spatial patterns, Splachnum, 161
Sphaerophorus melanocarpus, 216
Sphaerotheciella sphaerocarpa, 398
Sphagnum, 7+; checklist of species of North America north of Mexico, 500; sect. **Inretorta** sect. nov., 283; angustifolium, 17, 349; austinii, 17+; capillifolium, 17+; centrale, 17+; fuscum, 17+; **inretortum** sp. nov., 283*; jensenii, 17; lindbergii, 17; magellanicum, 17+, 349; majus, 17+; pacificum, 17+; palustre, 17+; papillosum, 17+; rubellum, 17+; russowii, 17+; squarrosum, 17+; tenellum, 17+; teres, 17+; warnstorfii, 17+
Spiridentopsis longissima, 400
Splachnum, spatial patterns and sex ratios in dioecious and monoecious species, 161; ampullaceum, 162+; rubrum, 162+; sphaericum, 162+

Sporastatia testudinea, 204
Squamarina lentigera, 204
Squamidium biforme, 401; leucotrichum, 401
Staurothele clopima, 204; fissa, 365
Stenocybe, 217; major, 217; minutissima, 217+
Stereocaulon commixtum, 155; dactylophyllum var. occidentale, 217; dendroides, 155; exutum, 156; intermedium, 156; japonicum, 156; nigrum, 156; octomerum, 156; ramulosum, *216*; rivulorum, 204; sorediiferum, 157; tomentosum, 204; verruculigrum, 157; vesuvianum, 157
Sticta fuliginosa, 266; nylanderiana, 120; weigelii, 266; yatabeana, 120
Stirtonia ramosa, new depsides in, 279
STONE, D. F. & B. MCCUNE. Annual branching in the lichen Evernia prunastri in Oregon, 32
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—121, 69; —121, 222; —123, 502
Strangospora moriformis, 364; pinicola, 364
Sulcaria sulcata, 148
Symphysodon neckeroides, 397+*; vitianus, 400
Symphysodontella attenuatula, 400; cylindracea, 400; parvifolia, 400
Synalissa melambola, *217+*
Syrrhopodon aberrans, *444*; albidus, *444*; apertifolius, *440*; boulayanum, *440*; brevicaulis, *440*; cataractarum, *444*; chlorosus, *440*; glaucophyllus, *440*, var. rufus, *440*; hispidocostatus, *443*; integrifolius, *440*; isleanus, *440*; kilaueae, *440*; leucophanoides, *440*; ligulifolius, *444*; littoralis, *440*; patulifolius, *440*; prolifer, 438+, var. acanthoneuros, 440*, var. **albidus** comb. nov., 444*, var. cincinnatus, 444+, var. **hispidocostatus** comb. nov., 443*, var. mossmanensis, 442*, var. papillosus, 442*, var. prolifer, 440+*, var. tenuifolius, 440+*, var. **tosaensis** comb. nov., 442*; spathulifolium, *440*; spiralis, *440*; tosaensis, *442*

Taiwanobryum speciosum, 399
TAN, B. C. Six new taxa of Malesian mosses, 429
Tephromela armeniaca, 204
Tetraphidopsis pusilla, 399
Tetraphis pellucida, 266
Thamnidiella longipes, *328*
Thamnium holzingeri, *330*; vancouveriense, *330*
Thamnobryum alopecurum, 401; plicatulum, 401; sandei, 401
Thamnolia subuliformis, 204; vermicularis, 104*, 204
Thelidium decipiens, 365; papulare, 365; parvulum, 217
Thelocarpon superellum, 365
Thelotrema abroselum, 217+; clandestinum, *217+*; lacteum, 217; leprocarpum, 217; platycarpum, 217
Thermutis velutina, 365
THIERS, B. M. [Review of:] H. Bischler-Causse. Marchantia L. The Asiatic and Oceanic taxa, 378
THOMSON, J. W. Ramalina unifolia sp. nov. from North America, 341
Thuidium delicatulum, 63, 266; minutulum, 266
TIMME, S. L. *See* BOWERS, F. D. & S. L. TIMME, 30
Tomenthypnum nitens, 17+, 62, 349
Toninia caeruleonigricans, 204; candida, 204; lobulata, 204, *217+*; squalida, 204
Trachyloma diversinerve, 400; indicum, 400+*; planifolium, 400
Trachylomataceae, 395+
Trachypodaceae, 395+
Trachypodopsis auriculata, 399; serrulata subsp. crispatula, 399
Trachypus bicolor, 399; humilis, 399
Trapelia placodioides, 365
Trapeliopsis, 217; flexuosa, 204
Treubia tasmanica, 409+*
Treubiales, 409+
Tricharia cretacea, 217
Tripterocladium brewerianum, 401
Tsuga canadensis, epiphytic lichens and bryophytes on, 257
TUCKER, S. C. [Review of:] M. E. Hale, Jr. & M. Cole. Lichens of California, 99
Tuckermannopsis oakesiana, 266+*, pinastri, 204
Tylophoron, 217; protrudens, 217

Ulota crispa, 266+
Umbilicaria deusta, 204; esculenta, 146; hirsuta, 204; hyperborea, 204, 425; kisovana, 146; krascheninnikovii, 204; vellea, 204, 389+, 424; virginis, 204
Usnea ceratina, 266; diffracta, 148; fulvoreagens, 204; hesperina, 266; hirta, 204; cf. nipparensis, 148; rubicunda, 266; strigosa, 220+; subfloridana, 266

Venezuela, Isopterygium acutifolium, new species in, 343
Verrucaria aethiobola, 365+; amylacea, 217; calciseda, 365; elaeomelaena, 217; funckii, 217, 365; fuscella, 366; sandstedei, 217
VITT, D. H. Schlotheimia brownii, 387*
VITT, D. H. *See* GIGNAC, L. D. & D. H. VITT, 7
VITT, D. H. & D. G. HORTON. Rich fen bryophytes in Missouri: Ecological comments and three state records, 62

WEBER, W. A. [Review of:] D. J. Galloway. Flora of New Zealand, Lichens, 382
Weymouthia billardieri, 401
WHITTEMORE, A. T. [Review of:] R. M. Schuster. The Hepaticae of South Greenland, 101
WILHELM, G. S. & J. R. BURKHALTER. Cladonia perforata, the northwest Florida population, 66
WONG, P. Y. & I. M. BRODO. Significant records from the lichen flora of southern Ontario, Canada, 357
WYATT, R. [Review of:] J. L. Doust & L. L. Doust (eds.). Plant reproductive ecology: Patterns and strategies, 382
WYATT, R. *See* CAMERON, R. G. & R. WYATT, 161

Xanthoparmelia chlorochroa, 204; coloradoensis, 217; coreana, 130; cumberlandia, 204; hypopsila, *217+*; idahoensis, 217; ioanis-simae, *217+*; kurokawae, *217+*; lavicola, 217+; lineola, 204, 319+*, 424; lipochlorochroa, 217; mexicana, 130, 204; monticola, 217+; neoconspersa, 217+; neowyomingica, 217; novomexicana, 204; piedmontensis, 131; planilobata, 217; plittii, 204, 424*; somloensis, 217+; subdecipiens, 204, 319+, 424; subramigera, 131; taractica, 204, 217+; wyomingica, 204
Xanthoria candelaria, 204; elegans, 204; fallax, 204; polycarpa, 52, 204; sorediata, 204, 366; subramulosa, 217

Yellowstone National Park, lichens of, 197

ZOMLEFER, W. B. & W. R. BUCK. A reassessment of four Rigodium types, 303